



Abbott Analytical



Consulting Scientists to the Disinfectant Industry

Certificate of Analysis

Sample(s) : One sample of Hygiene4less 70% Alcohol Hand Gel

Received from: Maclin Sourcing Solutions Ltd. Unit A3, Risby Business Park,
Newmarket Road, Risby, Bury St Edmunds, IP28 6RD

Date received: 12 October 2012 **Date tested:** 23 October 2012

Certificate no: 12K.076HR.MSS **Certificate date:** 24 October 2012

Sample ref: 12K/076 **Page:** 1 of 7

Analysis required: EN 1500, Chemical disinfectants and antiseptics - Hygienic handrub - Test method and requirements (phase 2, step 2)

Volume of product used: 6ml (2x 3ml)

Total rubbing time with product: 1 minute (2x 30 seconds)

Identification of bacterial strain(s) used: *Escherichia coli* (K12) NCIMB 10083

Number in contamination fluid (N): 2.85×10^8

Neutralising solution: 30g/l Polysorbate 80, 3g/l Lecithin,
1g/l Histidine, 1g/l Cysteine

Principle of test:

The number of test organisms released from the fingertips of artificially contaminated hands is assessed before and after using the hygienic handrub. The ratio of the two resulting values is called the reduction factor; it represents a measure of antimicrobial activity of the hygienic handrub product tested.

In order to achieve the necessary precision a large number of subjects has to be used because of the possible variation in bacterial flora found on human skin. In this case a total of 15 subjects (healthy adults) were chosen, each one carrying out the test procedure in precisely the same way as the others. To compensate for extraneous influences the test sample reduction factor (P) is compared with the reduction factor obtained with a parallel reference handrub procedure (R) which is performed with the same subjects, on the same day and under comparable environmental conditions.

D C Watson



Abbott Analytical



Consulting Scientists to the Disinfectant Industry

24 October 2012

Certificate No: 12K.076HR.MSS

Page 2 of 7

Experimental procedure:

1) Application of the contamination fluid

Each of the 15 subjects was asked to wash their hands for 1 minute in soft soap to remove natural commensal organisms and then dry them thoroughly on a paper towel. The hands were then contaminated with very large numbers of bacteria well in excess of that experienced in normal everyday conditions. The hands were immersed in the contamination fluid (containing an overnight culture of the test organism, in this instance *E. coli* (K12), at a concentration of approximately 10^8 cfu/ml) in a suitable sized container for 5 seconds. The hands were removed from the contamination fluid and surplus liquid allowed to drain back into the container. This time the hands were allowed to air dry for approximately 3 minutes holding them horizontally with fingers spread out and rotating them to and fro to avoid the formation of droplets.

2) Prevalues

Immediately after drying, each of the 15 subjects was asked to rub their fingertips, including the thumbs, for 1 minute on the base of a petri dish (a separate dish for each hand) containing 10ml of maximum recovery diluent (MRD) without neutraliser, in order to assess the release of test organisms before treatment of the hands. Dilutions of these sample fluids were prepared to 10^{-4} . A 1ml aliquot of each dilution for each hand was placed in a separate petri dish, 10-15ml of Tryptone Soy Agar (sterilised and cooled to 45°C) added and mixed thoroughly. Plates were allowed to set and incubated at 37°C for 24 hours. Each plate was then examined for growth of the test organism.

3) Hygienic handrub procedure

Each of the 15 subjects was asked to pour 3ml of 60% propan-2-ol into cupped hands and rub vigorously for 30s onto the skin, up to the wrists in accordance with the standard handrub procedure. This comprises five strokes backwards and forwards palm to palm, right palm over left dorsum and left palm over right dorsum, palm to palm with fingers interlaced, back of fingers to opposing palms with fingers interlocked, rotational rubbing of right thumb clasped in left palm and left thumb clasped in right palm, rotational rubbing with clasped fingers of right hand in palm of left hand and clasped fingers of left hand in right palm. Repeated with a further 3ml propan-2-ol to give a total rubbing time of 60 seconds. After 60 seconds the hands are rinsed under running tap water for 5 seconds, excess water is shaken off.

D C Watson



Abbott Analytical



Consulting Scientists to the Disinfectant Industry

24 October 2012

Certificate No: 12K.076HR.MSS

Page 3 of 7

4) Handrub procedure with test product (P)

The above procedure was repeated exactly using the test product in place of propan-2-ol. The volume of test product used and the total rubbing time were as detailed in the table on the first page.

5) Postvalues

Immediately after rinsing the 15 subjects were asked to rub the fingertips on the base of a petri dish containing 10ml of MRD with neutraliser for 1 minute using a separate petri dish for each hand. Then 1ml of each of the undiluted sample fluids was placed in a petri dish and covered with 15ml of TSA mixed thoroughly and allowed to set. Plates were then incubated overnight at 37°C and examined for growth of the test organism.

6) Calculation

The number of colony forming units (cfu) per plate for each dilution was recorded and the number of cfu/ml of sample fluid calculated. For both reference and test procedure the log counts from right and left hands of each subject were averaged separately for prevalues and postvalues.

From the difference between this individual combined log prevalue and the log postvalue a log reduction factor is established for each subject. Then the two arithmetic means of all individual log reduction factors are calculated for both the reference and the test procedure.

For the test product to pass the criteria of EN 1500 the mean log reduction factor obtained must not be significantly smaller than that obtained for the propan-2-ol.

Results: (see tables on following page)

See tables on following pages.

Difference of mean RFs: R-P = 5.53-5.60 = -0.07

Conclusion:

This batch of Hygiene4less 70% Alcohol Hand Gel [passes the requirements of EN 1500 for hygienic handrubs](#) when tested under the procedures described above.

D C Watson



Abbott Analytical



Consulting Scientists to the Disinfectant Industry

24 October 2012

Certificate No: 12K.076HR.MSS

Page 4 of 7

Results table: Reference handrub procedure

Subject		Number of cfu per plate from dilution 10 ^x				
Number	Hand	Prevalues		Postvalues		
		10 ⁻⁴	10 ⁻⁵	10 ⁰	10 ⁻¹	10 ⁻²
1	Left	244	28	8	2	0
	Right	288	35	11	2	0
2	Left	>300	36	9	0	0
	Right	284	30	15	2	0
3	Left	>300	35	18	3	0
	Right	>300	40	25	2	0
4	Left	272	30	20	2	0
	Right	268	30	14	0	0
5	Left	164	23	0	0	0
	Right	228	25	0	0	0
6	Left	288	35	21	3	0
	Right	>300	36	35	3	0
7	Left	>300	30	20	3	0
	Right	264	28	15	2	0
8	Left	252	28	0	0	0
	Right	300	31	0	0	0
9	Left	244	27	8	0	0
	Right	236	25	11	0	0
10	Left	218	19	0	0	0
	Right	226	23	0	0	0
11	Left	>300	40	15	2	0
	Right	>300	35	18	3	0
12	Left	>300	37	27	3	0
	Right	268	29	34	5	0
13	Left	184	20	0	0	0
	Right	222	23	0	0	0
14	Left	>300	29	38	5	0
	Right	278	27	51	5	1
15	Left	>300	41	14	2	0
	Right	>300	37	23	2	0

Numbers in **blue** used for further computation

D C Watson

D C Watson BSc, CBiol, MSB, MIFST, ACIEHO
PO Box 95, New Ferry, Wirral, CH62 6HA
Tel: 0151 637 3331 Mob: 07767 871275
email: abbottanalytical@hotmail.co.uk



Abbott Analytical



Consulting Scientists to the Disinfectant Industry

24 October 2012

Certificate No: 12K.076HR.MSS

Page 5 of 7

Results table: Handrub procedure with test product

Subject		Number of cfu per plate from dilution 10 ^x				
Number	Hand	Prevalues		Postvalues		
		10 ⁻⁴	10 ⁻⁵	10 ⁰	10 ⁻¹	10 ⁻²
1	Left	222	24	4	0	0
	Right	250	21	7	0	0
2	Left	>300	35	11	2	0
	Right	>300	38	15	2	0
3	Left	296	33	15	2	0
	Right	>300	35	21	2	0
4	Left	>300	27	10	0	0
	Right	252	23	14	0	0
5	Left	250	27	0	0	0
	Right	204	23	0	0	0
6	Left	>300	33	46	6	1
	Right	>300	36	55	5	1
7	Left	208	25	0	0	0
	Right	242	28	0	0	0
8	Left	216	23	0	0	0
	Right	267	31	0	0	0
9	Left	>300	35	15	2	0
	Right	>300	29	18	2	0
10	Left	208	24	5	0	0
	Right	216	25	3	0	0
11	Left	252	29	7	0	0
	Right	260	33	5	0	0
12	Left	>300	33	18	3	0
	Right	300	27	25	3	0
13	Left	256	18	0	0	0
	Right	204	17	0	0	0
14	Left	>300	38	31	4	0
	Right	>300	34	38	5	0
15	Left	>300	36	22	3	0
	Right	>300	29	14	1	0

Numbers in **blue** used for further computation

D C Watson

D C Watson BSc, CBiol, MSB, MIFST, ACIEHO
PO Box 95, New Ferry, Wirral, CH62 6HA
Tel: 0151 637 3331 Mob: 07767 871275
email: abbottanalytical@hotmail.co.uk



Abbott Analytical



Consulting Scientists to the Disinfectant Industry

24 October 2012

Certificate No: 12K.076HR.MSS

Page 6 of 7

Results table: Computed log values (means of left and right hand) and log reduction factors

Subject	Reference disinfection procedure (R)			Disinfection procedure with test product (P)		
	log x	log y	log z	log x	log y	log z
1	6.43	0.97	5.46	6.37	0.72	5.65
2	6.51	1.07	5.44	6.56	1.11	5.45
3	6.57	1.33	5.24	6.51	1.25	5.26
4	6.44	1.22	5.22	6.41	1.07	5.34
5	6.30	0.00	6.30	6.36	0.00	6.36
6	6.51	1.43	5.08	6.54	1.70	4.84
7	6.45	1.24	5.21	6.36	0.00	6.36
8	6.44	0.00	6.44	6.38	0.00	6.38
9	6.38	0.97	5.41	6.50	1.22	5.28
10	6.34	0.00	6.34	6.33	0.59	5.74
11	6.57	1.22	5.35	6.42	0.77	5.65
12	6.50	1.48	5.02	6.50	1.33	5.17
13	6.31	0.00	6.31	6.35	0.00	6.35
14	6.45	1.64	4.81	6.56	1.54	5.02
15	6.59	1.25	5.34	6.51	1.24	5.27
\bar{x}	6.45	0.92	5.53	6.44	0.84	5.60
s	0.09	0.60	0.54	0.08	0.60	0.53
N	15	15	15	15	15	15

log x: log prevalue

log y: log postvalue

log z: log reduction factor

\bar{x} : overall mean of log x, log y, log z

s: standard deviation

N: number of values (subjects) in each column

D C Watson

D C Watson BSc, CBiol, MSB, MIFST, ACIEHO
PO Box 95, New Ferry, Wirral, CH62 6HA
Tel: 0151 637 3331 Mob: 07767 871275
email: abbottanalytical@hotmail.co.uk



Abbott Analytical



Consulting Scientists to the Disinfectant Industry

24 October 2012

Certificate No: 12K.076HR.MSS

Page 7 of 7

Neutralization & toxicity validation

Test Organism	<i>Escherichia coli</i> (K12)	
Validation Suspension (N _v)	Vc1 234	Vc2 216
	$\bar{x} = 225$	
Neutraliser Control (B)	Vc1 228	Vc2 202
	$\bar{x} = 215 \geq 0.5N_{v_0}$	
Method Validation (C)	Vc1 206	Vc2 241
	$\bar{x} = 224 \geq 0.5N_{v_0}$	
Test Suspension	10 ⁻⁶ Vc1 258	Vc2 310
	10 ⁻⁷ Vc1 27	Vc2 33
(N)	$\bar{w} = 2.85 \times 10^8$	

Vc = plate count per ml

\bar{x} = average of Vc1 and Vc2

\bar{w} = weighted mean of \bar{x}

R = reduction ($\lg R = \lg N_0 - \lg N_a$)

D C Watson

D C Watson BSc, CBiol, MSB, MIFST, ACIEHO
PO Box 95, New Ferry, Wirral, CH62 6HA
Tel: 0151 637 3331 Mob: 07767 871275
email: abbottanalytical@hotmail.co.uk